

Mg-Al granulite garnet-orthopyroxene geothermometer gives temperatures between 700 and 840°C. Garnet-sillimanite-plagioclase-quartz geobarometer in a pelitic schist gives pressure of 2.5 kbar pressure at the garnet-biotite temperature of 620°C for the eastern part of the area. Pressure calculated using garnet-orthopyroxene-plagioclase-quartz geobarometer in a mafic granulite in the western

part of the area is 4.6 kbar at the garnet-orthopyroxene temperature of 790°C.

Acknowledgements: SST acknowledges CSIR, New Delhi for a research fellowship (No. 2-44/2000(i)-EU.II). Sri Sagnik Dasgupta of IIT, Bombay is thanked for contribution in structural mapping and interpretation. We thankfully acknowledge critical comments from two anonymous reviewers.

References

- BERMAN, R.G. (1991) Thermobarometry using multiequilibrium calculations: a new technique with petrological applications. *Can. Mineral.*, v.29, pp.833-855.
- CROOKSHANK, H. (1963) Geology of southern Bastar and Jeypore from the Bailadila Range to the Eastern Ghats. *Mem. Geol. Surv. India*, 87, 149 p.
- GHOSH, J.G. (2004) 3.56 Ga tonalite in the central part of the Bastar craton, India: oldest Indian date. *Jour. Asian Earth Sci.*, v.23, pp.359-364.
- GHOSH, P.K. (1941) The charnockite series of Bastar state and western Jeypore. *Rec. Geol. Surv. India*, v.91, Prof. Paper No.15, 55p.
- MISHRA, V.P., SINGH, P. and DUTTA, N.K. (1988) Stratigraphy, structure and metamorphic history of Bastar craton. *Rec. Geol. Surv. India*, v.117 (Parts 3-8), pp.1-26.
- RAMACHANDRA, H.M., ROY, A., MISHRA, V.P. and DUTTA, N.K. (2001) A critical review of the tectonothermal evolution of the Bastar craton. *Geol. Surv. India Spl. Publ. No.55 (Vol. II)*, pp.161-180.
- RAMAKRISHNAN, M. (1990) Crustal development in southern Bastar Central Indian craton. *Geol. Surv. India Spl. Publ. No.28*, pp.44-66.
- SPEAR, F.S. (1995) *Metamorphic Phase Equilibria and Pressure-Temperature-Time Paths*. Mineralogical Society of America Monograph, 799p.
- THAKUR, S.S., PATEL, S.C. and DASGUPTA, S. (2006) Structural pattern in Sukma supracrustal rocks near Nagaras, southern Bastar Craton. *Jour. Geol. Soc. India*, v.68(5), pp.769-773.

(Received: 9 December 2005; Revised form accepted: 18 July 2006)

Announcement

LAKSHMESWAR RAMA RAO BRITH CENTENTARY RESEARCH GRANTS: 2007-2008

A sum of Rs.10,000/- shall be made available during the financial year 2007-2008 to young research scholars for carrying out laboratory and field work in the field of palaeontology and stratigraphy.

Those interested in seeking assistance are requested to apply to the Society furnishing full particulars of academic qualifications, synopsis of the research project they are engaged in and the extent of assistance required. The application should be supported by their Research Guide and the Head of the Institution and reach the Secretary, Geological Society of India, No.63, 12th Cross, Basappa Layout, Gavipuram PO, P.B. No.1922, Bangalore - 560 019. **Telefax:** 080-26613352; **Phone:** 080-22422943; 09886028190; 09900256793 (M); **Email:** gsocind@gmail.com on or before 31st March 2007.