Vol. 3, Nos. 3 & 4: pp. 357, 1961

## ON THE NOMENCLATURE OF CUT-LEAVED TECOMA

## J. K. MAHESHWARI

Central National Herbarium, Botanical Survey of India, Sibpore, Howrah

The genus Tecoma Juss. emend. HBK. (Bignoniaceae) is represented in India by a single species, T. stans (Linn.) Juss. ex HBK. Nov. Gen. (ed. 4) 3: 144, 1818, said to be a native of South America. It is now very common in Indian gardens and is established as an escape in some parts of the country (Konkan and North Kanara, ex Cooke; Dharwar, Hohenacker 784; Pachmarhi Hills, ex Witt; Asirgarh, M. P., J. F. Duthie 3868, J. K. Maheshwari ; Jubbulpur, M. P., R. S. Hole 228). The typical form has leaflets that are only toothed. but there is a variety with graceful foliage in which the leaflets are narrow and are cut almost to the midrib. This, a cut-leaved form of Tecoma stans, was named by Sweet (in Hort. Brit. ed. 1, 2: 284, 1826) Tecoma incisa, based on an unpublished Bignonia incisa (" of gardeners"). The name Tecoma incisa was originally published as a nomen subnudum and seems to have been validated with a description by Loudon (in Hort. Brit. 243, 1830). De Candolle (Prodr. 9: 224, 1845) examined Bertero's collection of this form from Guadeloupe in the West Indies and renamed it Tecoma stans var. apiifolia DC. This name has been adopted by Benthall (in Trees Calcutta 336, 1946). According to the Rules of Nomenclature, the earliest validly published name

for this plant is *Tecoma incisa* Sweet ex Loudon. The specific name *incisa* must be reinstated as a varietal epithet under *Tecoma stans*. The correct name of the cut-leaved form is, therefore, as follows:

Tecoma stans Juss. ex HBK. var. incisa (Sweet) Maheshwari, comb. nov. Basi. Tecoma incisa Sweet, Hort. Brit. ed. 1, 2: 284, 1826, nomen subnudum; Loudon, Hort. Brit. 243, 1830, non Johnston (1940).
Syn. Tecoma stans var. apiifolia DC. Prodr. 9: 224, 1845; Benthall, Trees Calc. 336, 1946. Stenolobium stans D. Don var. apiifolium Seem. Jour. Bot. Lond. 1: 88, 1863; van Steenis, Rec. Trav. bot. néerl. 24: 906, 1927. Bignonia incisa Hortul.

Common name: Cut-leaved Tecoma.

An evergreen, ornamental shrub, about 3 m. high. Leaflets narrow,  $4-10 \times 1.5-2.5$  cm., deeply cut to the midrib and almost pinnatifid. Flowers yellow.

A native of Tropical America from Argentina to Mexico and now widely distributed and cultivated in the tropics. According to Koorders (Exc. Fl. Java 3: 184, 1912), it is in Java very frequently found in parks and gardens as well as in India, Burma and Malaya (ex Herb. CAL et SING).