NEW COMBINATION ON SOME VARIETIES OF *ELAEAGNUS UMBELLATA* SUBSP. *PARVIFOLIA* (WALL. EX ROYLE) SERVETTAZ

During the revisionary studies on the genus Elaeagnus it was found that E. umbellata Thunb. (1784) and E. parvifolia Wall. ex Royle (1836) are two distinct species. Schlechtendal (1857) also maintained this view but Servettaz (1908) treated E. parvifolia as a subsp. of E. umbellata and described some varieties under it. Momiyama & Hara (1973) have clearly stated that the Himalayan species E. parvifolia differs from E. umbellata on these points as stated below:

Buds and leaves with silvery, stellate scales. Perianth limb narrow, cylindrical with only silvery scales; lobes pointed and 2/3 of limb. Anthers with connectives. Styles short, copiously stellate pilose. Disc not hairy

Elaeagnus parvifolia Wall. ex Royle

Buds and leaves with brownish scales. Perianth limb broad conical with brown hairs; lobes 1/3 of limbs, rounded apex. Anthers sessile, pointed at apex. Styles long recurved. Disc hairy

E. umbellata Thunb.

Following this some new combinations have become necessary and proposed here.

Elaeagnus parvifolia Wall. ex Royle var. nuda (Serv.) Basu comb. nov.

Elaeagnus umbellata subsp. parvifolia (Wall. ex Royle) Serv. var. nuda Serv. in Beihe. Bot. Centralbl. (Abt. 2) 25: 57. 1909.

E. parvifolia Wall. ex Royle var. villosa (Serv.) Basu comb. nov.

E. umbellata subsp, parvifolia (Wall. ex Royle) Serv. var. villosa Serv. l.c. 57.

ACKNOWLEDGEMENTS

The author is thankful to the Director, Botanical Survey of India for providing all the facilities during the work and to the Deputy Director, Botanical Survey of India, Northern Circle, for going through the manuscript.

D. BASU

Botanical Survey of India, Dehra Dun

REFERENCES

Момічама, Ү. and H. Hara. Journ. Jap. Bot. 48(9): 266. Japan. 1973.

ROYLE, J. P. Illustrations of Botany and the Himalayan Mountains. 323, t. 81. f. 1. 1836.

Schlechtendal, F. L. In De Candolle: Prodromus 14:612.1857.

Servettaz, C. M. Bulletin De L' Herbier Boissier (2nd Serv.) 8(6): 383. 1908.

Beihefte zum Botanischen Centralblat (Abt. 2) 25: 56-57. Dresden. 1909.

THUNBERG, C. P. Flora Japonica 66, t. 14. 1784 (ed. 1975).

SOME NEW RECORDS OF THE GENUS MALVA L. (MALVACEAE) FROM INDIA

This paper records the occurrence of three taxa of the genus Malva L. from India for the first time. The specimens of these taxa were collected from north-western India and Rajasthan, deposited in the Central National Herbarium (CAL). Detailed analysis of these specimens showed that these are Malva mohileviensis Downar, M. verticillata L.

var. rafigii S. Abedin and M. parviflora L. var. microcarpa (Pers.) Loscos.

A detailed description of these taxa are given below to facilitate easy identification.

1. Malva mohileviensis Downar in Bull. Soc. Mosc. 1: 177. 1861; Iljin in Shish. & Borb., Fl. URSS 15: 64. Pl. 3 f. 1. 1949; S. Abedin, Fl. W. Pak. 130: 43. 1979.